

Basic Biostatistics

Looking beyond numbers

Jamalludin Ab Rahman MD MPH

Associate Professor

Department of Community Medicine

Kulliyyah of Medicine



Content

1. Basic premises – variables, level of measurements, probability distribution
2. Descriptive statistics
3. Inferential statistics, hypothesis testing







*We observe, we believe.
What we observe might not be the truth*



...and we can't observe all. We sample.



Population ➔ *Sample*


Parameter ➔ *Statistic*




Variable

- Characteristic of a population
- Can take different values
- Data = *measurements* collected/observed





Normal distribution



Characteristics *of Normal distribution*

- Smooth, symmetrical (around the mean), uni-modal, bell shaped curve
- Mean = Median = Mode
- Skewness = 0
- Kurtosis = 0
- The total area under the curve (AUC) = 1
- Asymptotic to the x-axis – never touch x-axis



Test of Normality

- Anderson–darling Test
- Corrected Kolmogorov–Smirnov Test (Lilliefors Test)
- Cramér–von-mises Criterion
- D'agostino's K-squared Test
- Jarque–Bera Test
- Pearson's Chi-square Test
- Shapiro–Francia
- Shapiro–Wilk Test




Use Normality test with caution

- Small samples almost always pass a normality test. *Normality tests have little power to tell whether or not a small sample of data comes from a Gaussian distribution.*
- With large samples, minor deviations from normality may be flagged as statistically significant, even though small deviations from a normal distribution won't affect the results of a t test or ANOVA.




Descriptive statistics



Association

- a value and whose associated value may be changed




e.g. Smoking


e.g. Lung Cancer




Disease model



Disease model (example)



Causal relationship



Hypothesis Testing

- Involve more than one variables
 - *exposure & outcome,*
 - *predictor & criterion,*
 - *risk & disease*
- Try to prove that
Exposure causes the Disease
e.g. Smoking causing Lung Cancer
- Example ~ H_0 : No difference of risk to get Lung Cancer between smoker and non-smoker



	Lung Cancer	No Lung Cancer
Smoking	20 (18.2%)	90 (81.8%)
Not Smoking	5 (4.5%)	105 (95.5%)

$$\chi^2 (df=1) = 10.150, p = 0.001, OR = 4.7 (CI95\% 1.7 - 13.0)$$

Because $p < 0.05$, we reject H_0 . Therefore there is a difference between smoker & non smoker



Statistical Test

- **Univariate** ~ One dependent & one independent
- **Multivariate** ~ Multiple dependent & multiple independent variable




What test to use?

Variable 1	Variable 2	Test
Categorical	Categorical	Chi-square
Categorical (2 pop)	Numerical (Normal)	Independent sample t-test
Categorical (2 pop)	Numerical (Not Normal)	Mann-Whitney U test
Categorical (> 2 pop)	Numerical (Normal)	One-way ANOVA
Categorical (> 2 pop)	Numerical (Not Normal)	Kruskal-Wallis test
Numerical (Normal)	Numerical (Normal)	Pearson Correlation Coefficient Test
Numerical (Normal/ Not Normal)	Numerical (Not Normal)	Spearman Correlation Coefficient Test
Numerical (Normal)	Numerical (Normal) – Paired	Paired t-test
Numerical (Not Normal)	Numerical (Not Normal) – Paired	Friedman test



But life is not simple!



Multivariate Analysis

- Hypothesis testing & control for confounders
 - *e.g. General Linear Model, Logistic Regression*
- Modeling
 - *e.g. Linear Regression*
- Data reduction
 - *e.g. Factor Analysis, Cluster Analysis*



Writing plan for statistical analysis

#1

Data were analyzed using the complex sample function of SPSS (version 13.0). Sampling errors were estimated using the primary sampling units and strata provided in the data set. Sampling weights were used to adjust for nonresponse bias and the oversampling of blacks, Mexican Americans, and the elderly in NHANES. The prevalence of hypertension, as well as the awareness, treatment, and control rates, were age adjusted by direct standardization to the US 2000 standard population.¹⁰ To analyze differences over time, the 2003–2004 data were compared with the 1999–2000 data. Estimates with a coefficient of variation >0.3 were considered unreliable. A 2-tailed P value <0.05 was considered statistically significant.

(Ong et al. 2009)



Writing plan for statistical analysis

#2

To assess the effect of the selection process on the characteristics of the cases, we compared cases included in the final analysis to the rest of the cases. Since controls included in the present analysis were different from the rest of the diabetes free participants by design, no similar comparisons were performed for that group. To compare baseline characteristics of cases and controls appropriate univariate statistics were used. Similar binary logistic and multiple linear regression models were built with incident diabetes or HbA1c as respective outcomes and additive block entry of adiponectin and potential confounders. For linear regression CRP and triglycerides were log transformed. Since HbA1c could be modified by drug treatment, we ran a sensitivity analysis excluding all participants on antidiabetic medication. A p-value of <0.05 was considered significant. Analyses were performed with SPSS 14.0 for Windows.



Reporting analysis (example)

TABLE 1. Characteristics of the cohort

	No known diabetes	Known diabetes	Total	P value
Admissions	62.72 (710)	37.28 (422)	1132	
Patients	64.78 (629)	35.22 (342)	971	
HbA1c (%)	6.05 ± 0.87	8.49 ± 2.56	6.96 ± 2.08	<0.001
HbA1c ≥7.0 (yes)	9.44 (67)	69.43 (293)	31.80 (360)	<0.001
Admission glucose (mg/dl)	118.39 ± 52.65	220.68 ± 175.32	156.52 ± 125.01	<0.001
Maximum glucose (mg/dl)	158.48 ± 87.85	318.98 ± 177.09	218.32 ± 150.13	<0.001
Glucose ≥200 mg/dl (yes)	17.61 (125)	72.04 (304)	37.90 (429)	<0.001
Age (yr)	56.62 ± 18.16	61.17 ± 14.70	58.32 ± 17.09	0.001
Sex (male)	50.70 (360)	47.63 (201)	49.56 (561)	0.40
Race/ethnicity				0.04
Black	27.32 (194)	30.57 (129)	28.53 (323)	
White	15.35 (109)	9.95 (42)	13.34 (151)	
Hispanic	41.97 (298)	41.23 (174)	41.70 (474)	
Other	15.35 (109)	18.25 (77)	16.43 (186)	
Prior medication/hospital/clinic (yes)	63.66 (452)	76.25 (321)	68.35 (773)	<0.001
HTN (yes)	53.10 (377)	87.68 (370)	65.99 (747)	<0.001
Systolic BP (mm/Hg)	136.51 ± 23.75	144.50 ± 25.93	139.50 ± 24.88	<0.001
Diastolic BP (mm/Hg)	76.43 ± 15.12	76.90 ± 14.68	76.61 ± 14.95	0.64
BMI (kg/m ²)	27.55 ± 7.62	28.88 ± 7.46	28.04 ± 7.58	0.03
LDL (mg/dl)	103.81 ± 42.81	98.04 ± 43.52	101.48 ± 43.16	0.13
HDL (mg/dl)	50.70 ± 20.88	45.45 ± 16.94	48.54 ± 19.52	<0.01
Triglycerides (mg/dl)	118.60 ± 89.33	161.19 ± 187.01	136.09 ± 139.56	<0.01

Data are presented as mean ± SD for continuous variables and percentage (number) for categorical variables based on total number of admissions. Totals may not equal 100% due to rounding. *P* values were calculated by generalized estimating equations. HTN, Hypertension; BP, blood pressure; LDL, low-density lipoprotein; HDL, high-density lipoprotein.



Reporting analysis (example)

Table 1. Sociodemographic Characteristics of the Participants

Sociodemographic Characteristics	Women (n=1800)			Men (n=1281)		
	n	Unweighted, %*	Weighted, %*	n	Unweighted, %*	Weighted, %*
Place of residence						
Urban	893	49.6	30.9	652	50.9	33.0
Rural	907	50.4	69.1	629	49.1	67.0
Age, y						
25 to 34	725	40.3	42.6	470	36.7	35.8
35 to 44	500	27.8	27.9	350	27.3	27.3
45 to 54	367	20.4	18.8	276	21.6	21.1
55 to 64	208	11.6	10.7	185	14.4	15.8
Education, y†						
None	685	38.1	44.7	196	15.3	21.2
1 to 4	642	35.7	37.3	423	33.1	36.1
5	191	10.6	7.5	248	19.4	19.4
6 to 7	164	9.1	6.1	199	15.6	12.2
≥8	115	6.4	4.4	213	16.6	11.2

*Within each variable, the sum of the proportions may not be 100% because of rounding.

†The sum of the number of participants in each category is <1800 for women and 1281 for men because of missing data.



Reporting analysis (example)

Table 2. Prevalence of Hypertension Among Women and Men From Urban and Rural Areas According to Age, Education, Body Mass Index, Waist Circumference, and Current Alcohol Drinking

Participant Characteristics	Hypertension							
	Women				Men			
	Urban		Rural		Urban		Rural	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI
All participants	41.0	36.4 to 45.7	26.8	19.5 to 34.1	40.0	34.0 to 46.1	33.5	27.8 to 39.2
Age, y								
25 to 34	17.6	12.7 to 22.5	11.1	5.9 to 16.2	31.8	25.0 to 38.6	32.8	26.4 to 39.2
35 to 44	43.2	31.8 to 54.5	27.1	18.8 to 35.5	35.3	25.7 to 44.9	27.7	18.8 to 36.6
45 to 54	69.5	57.6 to 81.3	45.5	35.6 to 55.4	49.8	39.6 to 60.1	32.0	20.3 to 43.7
55 to 64	73.0	64.0 to 81.9	57.9	44.9 to 70.9	59.4	38.3 to 80.5	46.0	46.0 to 60.8
Education, y								
0 to 4	45.8	38.0 to 53.6	27.0	19.7 to 34.3	42.7	26.3 to 59.1	33.7	26.1 to 41.3
5 to 7	37.6	32.4 to 42.9	18.5	7.6 to 29.4	39.0	28.3 to 49.7	32.8	21.8 to 43.8
≥8	27.2	20.8 to 33.6	46.8	4.6 to 89.0	38.3	29.5 to 47.2	35.0	4.0 to 66.0
Body mass index, kg/m ²								
<25.0	33.0	29.4 to 36.7	25.6	18.7 to 32.4	34.4	29.3 to 39.6	30.5	23.8 to 37.2
25.0 to 29.9	54.1	45.2 to 63.1	42.2	26.1 to 58.2	53.9	39.6 to 68.2	62.5	48.3 to 76.7
≥30	54.9	44.5 to 65.3	31.8	8.4 to 55.1	78.6	67.0 to 90.2	89.7	66.6 to 100.0
Waist circumference, cm								
Women <8 and men <102	38.0	33.2 to 42.8	27.4	19.9 to 34.9	38.3	32.5 to 44.1	32.5	27.0 to 38.0
Women ≥8 and men ≥102	60.0	50.4 to 69.6	50.9	26.1 to 75.7	79.3	65.3 to 93.2	100	*
Current drinking								
No	40.4	33.8 to 47.0	24.4	16.2 to 32.6	37.8	28.9 to 46.8	28.5	23.1 to 33.9
<1 d/wk	40.3	33.6 to 47.0	33.0	23.3 to 42.8	43.4	33.2 to 53.6	35.4	20.7 to 50.2
≥1 d/wk	48.2	33.1 to 63.3	34.6	20.1 to 49.1	38.6	27.7 to 49.5	40.5	31.4 to 49.6

*Only 1 subject was in this category.

